1992
Journal of Cardiopulmonary Rehabilitation
Subject Index

Age, and cardiac disease in public safety officers, 51–54
Angioplasty, coronary, return to work after, 20–24
Anticoagulants, oral, improving safety of, 15–17
Arkansas physicians, views on rehabilitation, 32–35
Arm exercise
on air-braked ergometer, cardiorespiratory responses to, 244–252
in chronic obstructive pulmonary disease, hemodynamic and respiratory responses to, D4
ergometry in prediction of treadmill performance, 36–41
postinfarction, with leg ergometry, 174–181
Asthma risk, with inhaled beta-agonists, 233–235
Atherosclerosis progression, life style changes affecting, 159–163

Beta-agonists, inhaled, and risk of fatal asthma, 233–235
Beta-blockers, and heart rate response to exercise, 202–206
Blood pressure
calcium intake affecting, 13–14
diet and exercise program in hypertension with diabetes, 194–200
measurements at rest and during exercise, 277–285
in prolonged walking, caffeine affecting, D3
Bypass surgery
effects on smoking behavior, D6
exercise testing after, prediction of oxygen uptake in, 111–115

Caffeine, and heart rate and blood pressure response to prolonged walking, D3
Calcium intake, affecting blood pressure, 13–14
Cardiopulmonary resuscitation, prolonged, by conditioned patients with cardiac disease, 208–212
Children, actual dietary intake compared to recommendations, 92–93
Cholesterol levels
endurance training affecting, 42–49
oat cereals affecting, 164–166
omega-3 fatty acids affecting, 288–292
reduction as secondary prevention measure, 87–90
Competence and quality in rehabilitation, managing for, 237–242
Compliance
and exercise endurance in chronic obstructive pulmonary disease, 105–110
with exercise programs, 18–19
with low-fat vegetarian diet in cardiac disease, D5
with outpatient rehabilitation services, 25–30
Coronary angioplasty, return to work after, 20–24
Coronary artery bypass surgery
effects on smoking behavior, D6
exercise testing after, prediction of oxygen uptake in, 111–115
Coronary artery disease
functional classes affecting energy expenditure in daily activities, 56–61
infarction in. *See* Myocardial infarction and prediction of oxygen uptake after bypass surgery, 111–115
secondary prevention by lowering cholesterol, 87–90

Daily activities
and energy expenditure in coronary heart disease of different functional classes, 56–61
postinfarction leisure and social activity after rehabilitation program, 117–124
Depression, in chronic obstructive pulmonary disease, 261–268
Diabetes mellitus, hypertension with, diet and exercise program in, 194–200
Diet. *See* Nutrition

Editorials, 87–90, 159–163
Energy expenditure
with air-braked ergometer, 244–252
in daily activities of coronary heart disease patients of different functional classes, 56–61
in treadmill walking on soft platform belt, D2
Ergometry
arm performance in prediction of treadmill capacity, 36–41
postinfarction arm crank exercise with leg ergometry, 174–181
responses to arm ergometry exercises in chronic obstructive pulmonary disease, D4
responses to leg and arm work in air-braked ergometer, 244–252
Exercise
beta-blockers affecting heart rate in, 202–206
blood pressure measurements in, 277–285
caffeine affecting heart rate and blood pressure in prolonged walking, D3
gender differences in cardiac response to, 183–186
and life style changes affecting atherosclerosis progression, 159–163
walking with ankle and wrist weights in cardiac disease, 254–259
Exercise testing
after bypass surgery, and prediction of oxygen uptake, 111–115
arm ergometry and treadmill performance in, 36–41
energy expenditure in coronary heart disease of different functional classes, 56–61
oxygen cost and heart rate in treadmill walking on soft platform belt, D2
postinfarction arm crank with leg ergometry in, 174–181
responses to maximal treadmill and arm exercises in chronic obstructive pulmonary disease, D4
Exercise training
benefits in chronic obstructive pulmonary disease, 188–192
cholesterol levels in, 42–49
compliance affecting endurance in chronic obstructive pulmonary disease, 105–110
dietary program with, in diabetes with hypertension, 194–200
evolution at San Diego State University, 70–71
improving compliance with, 18–19
and prolonged cardiopulmonary resuscitation by conditioned patients, 208–212
responses to leg and arm work in air-braked ergometer, 244–252
and ventilatory threshold in heart transplant recipients, 126–129
walking with ankle and wrist weights in cardiac disease, 254–259

Fatty acids in n-3 family, affecting cardiovascular health, 288–292
Fish oil intake, affecting cardiovascular health, 288–292

Gender, and left ventricular response to exercise, 183–186

Heart disease
compliance with low-fat vegetarian diet in, D5
prolonged CPR by conditioned patients in, 208–212
walking with ankle and wrist weights in, 254–259
Heart rate
comparison of two pacemaker models in patients with pulmonary disease, 270–275
in exercise, beta-blockers affecting, 202–206
in prolonged walking, caffeine affecting, D3
Heart transplantation, and ventilatory threshold after exercise training, 126–129
Hemodynamic responses, in maximal treadmill and arm exercises in chronic obstructive pulmonary disease, D4
Hypertension in diabetes mellitus, diet and exercise program in, 194–200

Infarction. *See* Myocardial infarction

Leg exercise
on air-braked ergometer, cardiorespiratory responses to, 244–252
postinfarction, arm crank exercise with, 174–181
Letters to the editor, 302–303
Life style changes, affecting atherosclerosis progression, 159–163
Lipid levels. *See* Cholesterol levels

Managed-care organizations, negotiations with, 170–171
Management practices. *See* Practice issues
Myocardial infarction
and leisure and social activities after rehabilitation program, 117–124
postinfarction arm crank with leg ergometry in exercise testing, 174–181
reinfarction prevention by lowering cholesterol, 87–90

Nitroglycerin, properties of, 94–97
Nutrition
actual intake of young children compared to recommendations, 92–93
calcium intake affecting blood pressure, 13–14
compliance with low-fat vegetarian diet in cardiac disease, D5
intensive diet and exercise program in diabetes with hypertension, 194–200
and life style changes affecting atherosclerosis progression, 159–163
lipid-lowering effects of oat cereals, 164–166
omega-3 fatty acids affecting cardiovascular health, 288–292

Oat cereals, lipid-lowering effects of, 164–166
Obstructive pulmonary disease, chronic (COPD)
benefits of exercise training in, 188–192
comparison of two adaptive pacemakers in, 270–275
depression in, 261–268
exercise compliance affecting endurance in, 105–110
responses to maximal treadmill and arm exercises in, D4
Occupational therapist, role of, 98–104
Omega-3 fatty acids, affecting cardiovascular health, 288–292
Outpatient rehabilitation services, attendance in, 25–30
Oxygen cost
in daily activities of coronary heart disease patients of different functional classes, 56–61
in treadmill walking on soft platform belt, D2
Oxygen uptake
prediction from treadmill performance after bypass surgery, 111–115
and ventilatory threshold after exercise training in heart transplant recipients, 126–129
in work on air-braked ergometer, 244–252

Pacemakers, comparison of two models in patients with pulmonary disease, 270–275
Practice issues
compliance with exercise programs, 18–19
costs of rehabilitation services, 171–173
managed-care negotiations, 170–171
operations management modeling to determine optimal case mix of three cardiac services, D-1
quality management methods, 237–242
role of occupational therapist, 98–104
Psychologic outcome
in depression with chronic obstructive pulmonary disease, 261–268
occupational therapy affecting, 98–104
Public safety officers, cardiac disease in, 51–54
Pulmonary patients. *See also* Obstructive pulmonary disease, chronic (COPD)
asthma risk with inhaled beta-agonists, 233–235

Quality management methods, 237–242
Quality of life
after coronary angioplasty, 20–24
occupational therapy affecting, 98–104
pulmonary rehabilitation program affecting, 105–110

Rehabilitation programs
affecting postinfarction leisure and social activities, 117–124
after coronary angioplasty, 20–24
attendance in outpatient services, 25–30
and bypass surgery affecting smoking behavior, D6
compliance affecting exercise endurance in chronic obstructive pulmonary disease, 105–110
cost of services in, 171–173
and depression in chronic obstructive pulmonary disease, 261–268
evolution at San Diego State University, 70–71
life style changes in, 159–163
occupational therapist in, 98–104
views of Arkansas physicians on, 32–35
Relaxation techniques, and life style changes affecting atherosclerosis progression, 159–163
Respiratory responses, in maximal treadmill and arm exercises in COPD, D4
Risk factors
evaluation in public safety officers, 51–54
serum cholesterol levels, 87–90

Safety of oral anticoagulants, improvement of, 15–17
San Diego State University, evolution of cardiac rehabilitation in, 70–71
Sex-related differences, in cardiac response to exercise, 183–186
Smoking behavior, coronary artery bypass surgery affecting, D6
Stress reduction, and life style changes affecting atherosclerosis progression, 159–163
Sudden death, and risk factors in public safety officers, 51–54

Theophylline, toxicity of, 167–169
Transplantation of heart, and ventilatory threshold after exercise training, 126–129
Treadmill exercise
after bypass surgery, prediction of oxygen uptake in, 111–115
in chronic obstructive pulmonary disease, hemodynamic and respiratory responses to, D4
performance prediction with arm ergometry, 36–41
on soft platform belt, oxygen cost and heart rate response in, D2

Ventilatory threshold, after exercise training in heart transplant recipients, 126–129

Walking
with ankle and wrist weights, in cardiac disease, 254–259
prolonged, caffeine affecting heart rate and blood pressure in, D3
Warfarin, improving safety of, 15–17
Weighted exercise, in walking with ankle and wrist weights in cardiac disease, 254–259
Work activity
after coronary angioplasty, 20–24
assessment by occupational therapist, 98–104
functional classes of coronary heart disease affecting energy expenditure in, 56–61
sudden cardiac death in, 51–54